

ESTABLISHED 2017

ELECTRIC CITY DENTAL

DR. VERONA SULJA

FAMILY DENTISTRY

DR. KOREN BENNETTS

PRO-SYNERGIX premium

TAKE-HOME WHITENING SYSTEM IN 3 EASY STEPS

STEP 1

OBJECTIVES

- TOOTH SENSITIVITY TREATMENT
- IMPROVEMENT OF TOOTH WHITENERS & GLOSS
- IMPROVEMENT OF ORAL HEALTH

YOUR TOOL

X-PUR Remin toothpaste

- 10% Nano Medical Hydroxyapatite ($mHAP$): to treat tooth sensitivity, improve tooth whiteness & gloss, remove stains and significantly improve your oral health.
- 10% Xylitol: refreshing taste
- Sodium Lauryl Sulfate & Triclosan free
- Lightly flavored

DIRECTIONS FOR USE

- 7 days prior to your whitening treatment (or during the production of your custom-made whitening trays supplied by your dental team)
 - Use as your daily toothpaste twice a day (morning & evening) before, during and after your treatment.
1. Place about 1 to 1.5 cm of X-PUR Remin on an ultra soft toothbrush and brush meticulously for 3 minutes.
 2. Expectorate. For best results, do not eat, drink or rinse for 30 minutes after brushing.

STEP 2

OBJECTIVE

- ULTRA-FAST WHITENING ACTION


YOUR TOOLS

- 2 x Purple syringes (1.2 ml each)
- Hydrogen Peroxide (H_2O_2)

DIRECTIONS FOR USE

Wear your trays once a day for 4 consecutive days or until there is no more gel.

For an even more accelerated whitening action
Wear your trays twice a day with newly dispensed gel. Allow twelve hours between applications.

1. Using a Purple syringe, dispense a small amount of gel* to the face-side of each tooth imprint in your custom-made trays. Each Purple syringe has calibration lines. Each line represents approximately 1 tray application.
* Example of the quantity of gel: 
2. Wear your trays for 30-60 minutes.
3. Rinse teeth and trays with water immediately following this step.

STEP 3

OBJECTIVE


- LONG-LASTING & ENHANCED WHITENING RESULTS

YOUR TOOLS

- 4 x Silver syringes (3 ml each)
- Carbamide Peroxide ($CH_3N_2O_3$)

DIRECTIONS FOR USE

Wear your trays once a day for 12 consecutive days or until desired shade is achieved.

1. Using a Silver syringe, dispense a small amount of gel* to the face-side of each tooth imprint in your custom-made trays. Each Silver syringe has calibration lines. Each line represents approximately 1 tray application.
* Example of the quantity of gel: 
2. Wear trays for 60-90 minutes.
For an even more accelerated whitening action
Wear your trays overnight.
3. Rinse teeth and trays with water immediately following this step.
4. If you are not satisfied with your whitening results, repeat Step 3 and contact your dental team for additional support.
5. Refrigerate any remaining product for future touch-ups, up to 12 months.

X-PUR Pro-Synergix Premium take-home whitening system is safe and effective when used according to the above directions for use. Your treatment must also be monitored by your dental team. This includes an initial diagnosis, professional teeth cleaning, review of directions and recall exam.

For more information, call Oral Science's Clients Service at 1 888 442.7070.



www.x-pur.com

ADVICE FOR USAGE

- Depending on your nutrition and lifestyle habits, whitening treatment may require a touch-up periodically (provided by your dental team).
- Reduce consumption of coffee, tea, red wine, cola and tobacco during your treatment, as they may contribute to tooth staining.
- Enamel, dentin, existing fillings and bonding materials are not affected or harmed by X-PUR Pro-Synergix Premium whitening agents. Allow 7-10 days to elapse following whitening, prior to any dental treatment requiring new bonding.

PRECAUTIONS

- Do not use whitening gel while pregnant or lactating.
- Keep whitening gel out of reach of children.
- Do not swallow whitening gel.
- Store syringes in a cool, dry place, or refrigerate. Do not freeze syringes.
- Do not expose syringes to heat or sunlight.

Enjoy your bright smile!

1 - 999 LANSDOWNE ST. W. PETERBOROUGH ON K9J 8N2

Tel: 705 749 9994 Fax: 705 742 8701 email: info@electriccitydental.ca www.electriccitydental.ca

Sulja Bennetts Dentistry Professional Corporation